Index to Volume 70
Author Index

Page number of erratum is in italic.

Abou-Arab, A. A. *See* H. A. Al-Kahtani, 619
Aboubacar, A. *See* A. A. Mohamed, 739
Al-Kahtani, H. A., and A. A. Abou-Arab. Comparison of physical, chemical, and functional properties of *Moringa peregrina* (Al-Yassar or Al-Ban) and soybean proteins, 619
Almeida-Dominguez, H. D., M. H. Gomez, S. O. Serna-Saldivar, R. D. Waniska, L. W. Rooney, and E. W. Lusas. Extrusion cooking of pearl millet for production of millet-cowpea weaning foods, 214
——, S. O. Serna-Saldivar, M. H. Gomez, and L. W. Rooney. Production and nutritional value of weaning foods from mixtures of pearl millet and cowpeas, 14
——. *See* S. O. Serna-Saldivar, 762
Andrews, J. L., M. J. Blundell, and J. H. Skerritt. A simple antibody-based test for dough strength. III. Further simplification and collaborative evaluation for wheat quality screening, 241
Aoe, S., T. Oda, K. Tatsumi, M. Yamauchi, and Y. Ayano. Extraction of soluble dietary fibers from defatted rice bran, 423
Ashworth, P. *See* S. L. Edward, 748
Ayano, Y. *See* S. Aoe, 423

Ballance, G. M. *See* A.-M. Bernier, 127
Balogh, A. *See* J. H. Wong, 113
Balshaw, R. *See* L. J. Malcolmson, 1, 417
Bason, M. L., J. A. Ronalds, C. W. Wrigley, and L. J. Hubbard. Testing for sprout damage in malting barley using the Rapid Visco-Analyser, 269
Bechtel, D. B., I. Zayas, R. Dempster, and J. D. Wilson. NOTE: Size-distribution of starch granules isolated from hard red winter and soft red winter wheats, 238
Bello, A. B. *See* A. G. W. Bradbury, 543
BeMiller, J. N. *See* J. E. Fannon, 611
Bennett, K. E., T. J. Siebenmorgen, and A. Mauromoustakos. Effects of McGill No. 2 miller settings on surface fat concentration of head rice, 734
Bernier, A.-M., and G. M. Ballance. Induction and secretion α-amylase, (1→3),(1→4)-β-glucanase, and (1→3)-β-glucanase activities in gibberellic acid and $CaCl_2$-treated half seeds and aleurones of wheat, 127
Bettge, A. D., and Y. Pomeranz. Air-aspirated cleaning to separate sound from preharvest-sprouted wheat, 36
Bhatty, R. S. Extraction and enrichment of (1→3),(1→4)-β-D-glucan from barley and oat brans, 73
——. Physicochemical properties of roller-milled barley bran and flour, 397
Bietz, J. A. *See* M. A. Dombrink-Kurtzman, 105
——. *See* F. R. Huebner, 506
Biliaderis, C. G., J. R. Tonogai, Perez, C. M., and B. O. Juliano. Thermophysical properties of milled rice starch as influenced by variety and parboiling method, 512
——. *See* N. M. Edwards, 122
——. *See* C. M. Perez, 567
Blundell, M. J. *See* J. L. Andrews, 241
Boros, D., R. R. Marquardt, B. A. Slominski, and W. Guenter. Extract viscosity as an indirect assay for water-soluble pentosan content in rye, 575
Boyer, C. D. *See* R. C. Yuan, 81
Bradbury, A. G. W., and A. B. Bello. Determination of molecular size distribution of starch and debranched starch by a single procedure using high-performance size-exclusion chromatography, 543
Breslauer, K. J. *See* G. Kaletunc, 548
Bruinsma, B. L. *See* K. Siffring, 351
Brusewitz, G. H. *See* X. Zhang, 133
Buchanan, B. B. *See* J. H. Wong, 113
Buriak, P. *See* S. R. Eckhoff, 723
Bushuk, W. *See* L. Gao, 475

Campaña, L. E., M. E. Sempé, and R. R. Filgueira. NOTE: Physical, chemical, and baking properties of wheat dried with microwave energy, 760
Campbell, G. M., C. D. Rielly, P. J. Fryer, and P. A. Sadd. Measurement and interpretation of dough densities, 517
Casasnovas, J. *See* G. R. Ziegler, 247
Champagne, E. T., and R. J. Hron, Sr. Utilizing ethanol containing an antioxidant or chelator to produce stable brown rice products, 562
Chan, K.-Y., and B. P. Wasserman. Direct colorimetric assay of free thiol groups and disulfide bonds in suspensions of solubilized and particulate cereal proteins, 22
——, and ——. Rapid solid-phase determination of cereal protein using bicinchoninic acid, 27
Chandrashekar, A. *See* H. Mazhar, 667
Chauhan, B. M. *See* A. Kumar, 504
Chen, Y. R. *See* N. Sharma, 59
Cheryan, M. *See* A. Mannheim, 115
Chinnaswamy, R. *See* A. S. Sokhey, 260
Chiu, M. M. *See* T. S. Kahlon, 435
Cho, K., and P. J. White. Enzymatic analysis of β-glucan content in different oat genotypes, 539
Chow, F. I. *See* T. S. Kahlon, 435
Chung, O. K. *See* G. L. Lookhart, 430
Cleemput, G., S. P. Roels, M. Van Oort, P. J. Grobet, and J. A. Delcour. Heterogeneity in the structure of water-soluble arabinoxylans in European wheat flours of variable bread-making quality, 324
——. *See* S. P. Roels, 318
——. *See* S. Vanhamel, 306
Cornejo, L., M. Hernández, and A. Sotelo. Nutritive value of chicken and corn flour mixtures in formulas for infants with lactose intolerance, 572
Cox, T. S. *See* G. L. Lookhart, 430
Crombez, M. *See* R. C. Eerlingen, 339
Cuevas-Perez, F. *See* B. O. Juliano, 650
Curran, S. P. *See* D. E. Rogers, 183
Czuchajowska, Z., and Y. Pomeranz. Protein concentrates and prime starch from wheat flours, 701

Dahlin, K. M., and K. J. Lorenz. Carbohydrate digestibility of laboratory-extruded cereal grains, 329
Darius, P. L. *See* S. Vanhamel, 306
Deceuninck, M. *See* R. C. Eerlingen, 345
D'Egidio, M. G., B. M. Mariani, S. Nardi, and P. Novaro. Visco-elastograph measures and total organic matter test: Suitability in evaluating textural characteristics of cooked pasta, 67
——, S. Nardi, and V. Vallega. Grain, flour, and dough characteristics of selected strains of diploid wheat, *Triticum monococcum* L., 298
——. *See* P. Novaro, 716
DeFloor, I., and J. A. Delcour. NOTE: Impact of milling procedure on breadmaking potential of cassava flour in wheatless breads, 616
——, M. Nys, and J. A. Delcour. Wheat starch, cassava starch, and cassava flour impairment of the breadmaking potential of wheat flour, 526
Delcour, J. A. *See* G. Cleemput, 324
——. *See* I. DeFloor, 526, 616
——. *See* R. C. Eerlingen, 339, 345
——. *See* S. P. Roels, 318
——. *See* S. Vanhamel, 306
——. *See* C. J. A. Vinkx, 311
Delpuech, B. *See* O. Neyreneuf, 109
Delwiche, S. R., and K. H. Norris. Classification of hard red wheat by near-infrared reflectance spectrocopy, 29
Dempster, R. *See* D. B. Bechtel, 238
Dexter, J. E. *See* N. M. Edwards, 122
——. *See* S. J. Symons, 90
Dick, J. W. *See* L. A. Grant, 676
Dimberg, L. H., O. Theander, and H. Lingnert. Avenanthramides—A group of phenolic antioxidants in oats, 637
Dombrink-Kurtzman, M. A., and J. A. Bietz. Zein composition in hard and soft endosperm of maize, 105
Dowd, M. E., P. J. Reilly, and W. S. Trahanovsky. Low molecular weight organic composition of ethanol stillage from corn, 204
Dubord, C. *See* P. Gélinas, 219
Dunkle, L. D. *See* B. P. Marks, 656

Eckhoff, S. R., W. V. Jayasena, and C. K. Spillman. Wet milling of maize grits, 257
——, K. D. Rausch, E. J. Fox, C. C. Tso, X. Wu, Z. Pan, and P. Buriak. A laboratory wet-milling procedure to increase reproducibility

and accuracy of product yields, 723
———. *See* E. J. Fox, 402
———. *See* A. H. Mistry, 360
———. *See* K. D. Rausch, 489
———. *See* P. J. Wu, 443
Edward, S. L., A. S. Hill, P. Ashworth, J. Matt, and J. H. Skerritt. Analysis of the grain-protectant pesticides chlorpyrifos-methyl and methoprene with a 15-min immunoassay for field or elevator use, 748
Edwards, N. M., M. S. Izydorczyk, J. E. Dexter, and C. G. Biliaderis. Cooked pasta texture: Comparison of dynamic viscoelastic properties to instrumental assessment of firmness, 122
Eerlingen, R. C., M. Crombez, and J. A. Delcour. Enzyme-resistant starch. I. Quantitative and qualitative influence of incubation time and temperature of autoclaved starch on resistant starch formation, 339
———, M. Deceuninck, and J. A. Delcour. Enzyme-resistant starch. II. Influence of amylose chain length on resistant starch formation, 345
Eggum, B. O., B. O. Juliano, C. M. Perez, and G. S. Khush. Starch, energy, and protein utilization by rats in milled rice of IR36-based amylose extender mutant, 275
Eisenbraun, G. J. *See* G. S. Ranhotra, 236
Eliasson, A.-C. *See* L. Wannerberger, 196
Ereifej, K. I., and R. Shibli. NOTE: Rheological properties and baking quality of flour from a landrace and durum wheat cultivars grown in Jordan, 486

Fannon, J. E., J. M. Shull, and J. N. BeMiller. NOTE: Interior channels of starch granules, 611
Faubion, J. M. *See* C. Kerr, 633
———. *See* D. M. Larsen, 647
Felker, F. C., and J. W. Paulis. Quantitative estimation of corn endosperm vitreosity by video image analysis, 685
Figueroa, J. D. C., and K. Khan. Albumin fraction from spring, winter, and soft wheats—protein and associated carbohydrate identified by gel filtration chromatography and gel electrophoresis, 689
Filgueira, R. R. *See* L. E. Campaña, 760
Fox, E. J., and S. R. Eckhoff. Wet milling of soft-endosperm, high-lysine corn using short steep times, 402
———. *See* S. R. Eckhoff, 723
Friend, C. P., R. D. Waniska, and L. W. Rooney. Effects of hydrocolloids on processing and qualities of wheat tortillas, 252
Fryer, P. J. *See* G. M. Campbell, 517
Fulcher, R. G. *See* S. S. Miller, 231

Gaines, C. S. *See* W. R. Windham, 662
Gao, L., and W. Bushuk. Polymeric glutenin of wheat lines with varying number of high molecular weight glutenin subunits, 475
Gélinas, P., M. Lagimonière, and C. Dubord. Baker's yeast sampling and frozen dough stability, 219
Gelroth, J. A. *See* G. S. Ranhotra, 236, 303, 364, 530
———. See D. E. Rogers, 558
Gennadios, A., C. L. Weller, and R. F. Testin. Modification of physical and barrier properties of edible wheat gluten-based films, 426
Ghaedian, A. R. *See* R. L. Wehling, 720
Gibson, T. S., C. J. Kaldor, and B. V. McCleary. Collaborative evaluation of an enzymatic starch damage assay kit and comparison with other methods, 47
Gidley, M. J. *See* W. R. Morrison, 385
Glaser, B. K. *See* G. S. Ranhotra, 303, 364, 530
Gomez, M. H. *See* H. D. Almeida-Dominguez, 14, 214
———. *See* R. Young, 179
Grant, D. L. *See* R. Graybosch, 95
Grant, L. A., J. W. Dick, and D. R. Shelton. The effects of drying temperature, starch damage, sprouting, and additives on spaghetti quality characteristics, 676
Graybosch, R., C. J. Peterson, K. J. Moore, M. Stearns, and D. L. Grant. Comparative effects of wheat flour protein, lipid, and pentosan composition in relation to baking and milling quality, 95
———, Y. W. Seo, and C. J. Peterson. Detection of wheat-rye chromosomal translocations using an antisecalin monoclonal antibody, 458
Grenus, K. M. *See* F. Hsieh, 493
Griffin, V. K. *See* B. R. Hamaker, 377
Grobet, P. J. *See* G. Cleemput, 324
———. *See* C. J. A. Vinkx, 311
Grosch, W. *See* R. Sarwin, 553
Gruchala, L., and Y. Pomeranz. Enzyme-resistant starch: Studies using differential scanning calorimetry, 163
Guenter, W. *See* D. Boros, 575

Hagiwara, S. *See* Y. Hibi, 671
Hamaker, B. R., and V. K. Griffin. Effect of disulfide bond-containing protein on rice starch gelatinization and pasting, 377
———. *See* A. A. Mohamed, 739
Hamann, D. D. *See* S. S. Heddleson, 744
Hanna, M. A. *See* N. Sharma, 59, 63
Hara, H. *See* K. Shiiba, 209
Hatcher, D. W., and J. E. Kruger. Distribution of polyphenol oxidase in flour millstreams of Canadian common wheat classes milled to three extraction rates, 51
———. *See* J. E. Kruger, 234
Hay, R. L. Effect of flour quality characteristics on puff pastry baking performance, 392
Heddleson, S. S., D. D. Hamann, and D. R. Lineback. The Dahlquist criterion: Applicability of a rheological criterion to the loss of pressure-sensitive tack in flour-water dough, 744
Hernández, M. *See* L. Cornejo, 572
Hibi, Y., T. Matsumoto, and S. Hagiwara. Effect of high pressure on the crystalline structure of various starch granules, 671
Hill, A. S. *See* S. L. Edward, 748
Hizukuri, S. *See* C. P. Villareal, 753
Hooper, D. G. *See* R. L. Wehling, 720
Hoseney, R. C. *See* C. Kerr, 633
———. *See* W. D. A. Lin, 448
———. *See* R. A. Miller, 585
———. *See* D. E. Rogers, 183
———. *See* K. A. Tilley, 602
Hron, R. J., Sr. *See* E. T. Champagne, 562
Hsieh, F., K. M. Grenus, L. Hu, and H. E. Huff. Twin-screw extrusion of rice flour with salt and sugar, 493
Hu, L. *See* F. Hsieh, 493
Huang, J. J., and P. J. White. Waxy corn starch: Monoglyceride interaction in a model system, 42
Hubbard, L. J. *See* M. L. Bason, 269
Huebner, F. R., and J. A. Bietz. Improved chromagraphic separation and characterization of ethanol-soluble wheat proteins, 506
Huff, H. E. *See* F. Hsieh, 493
Hussain, A., and O. M. Lukow. NOTE: Relation of D-zone omega gliadins to the proteins associated with differences in quality of durum wheats, 483

Ilangantileke, S. *See* V. Laksanalamai, 381
Islas-Rubio, A. *See* S. O. Serna-Saldivar, 762
Izydorczyk, M. S., and C. G. Biliaderis. Structural heterogenity of wheat endosperm arabinoxylans, 641
———. *See* N. M. Edwards, 122

Jackson, D. S. *See* R. L. Wehling, 720
Jane, J. *See* Y.-J. Wang, 171, 521
Jayasena, W. V. *See* S. R. Eckhoff, 257
Johnson, D. L. *See* G. S. Ranhotra, 303
Johnson, L. A. *See* C. C. Lee, 18
Juliano, B. O., C. M. Perez, and F. Cuevas-Perez. Screening for stable high head rice yields in rough rice, 650
———, and ———. NOTE: Critical moisture content for fissures in rough rice, 613
———. *See* C. G. Biliaderis, 512
———. *See* B. O. Eggum, 275
———. *See* C. M. Perez, 567
———. *See* A. P. Resurreccion, 101
———. *See* C. P. Villareal, 753

Kahlon, T. S., F. I. Chow, B. E. Knuckles, and M. M. Chiu. Cholesterol-lowering effects in hamsters of β-glucan-enriched barley fraction, dehulled whole barley, rice bran, and oat bran and their combinations, 435
Kaldor, C. J. *See* T. S. Gibson, 47
Kaldy, M. S., G. R. Kereliuk, and G. C. Kozub. Influence of gluten components and flour lipids on soft white wheat quality, 77
Kaletunc, G., and K. J. Breslauer. Glass transitions of extrudates: Relationship with processing-induced fragmentation and end-product attributes, 548
Kang, Y. Y. *See* K. O. Kim, 440
Kasarda, D. D. *See* S. Masci, 581
Keagy, P. M., and T. F. Schatzki. Machine recognition of weevil damage in wheat radiographs, 696
Kephart, K. D. *See* E. Souza, 280
Kereliuk, G. R. *See* M. S. Kaldy, 77
Kerr, C., R. C. Hoseney, and J. M. Faubion. Mixograph studies. VI. Combined effects of charge (pH), activated double-bond compounds, and oxidants on dough mixing properties, 633
Khan, K. *See* J. D. C. Figueroa, 689

Khush, G. S. *See* B. O. Eggum, 275
Kim, K. O., M. K. Kim, Y. Y. Kang, and Y. C. Lee. Effects of malting conditions on quality characteristics of malt and roasted malt extract, 440
Kim, M. K. *See* K. O. Kim, 440
Kim, W. S., and P. A. Seib. Apparent restriction of starch swelling in cooked noodles by lipids in some commercial wheat flours, 367
Kirleis, A. W. *See* J. J. Watterson, 452
Kiss, F. *See* J. H. Wong, 113
Klopfenstein, C. F. *See* W.-M. Wang, 707, 712
Knuckles, B. E. *See* T. S. Kahlon, 435
Knutson, C. A. Reaction patterns of maize α-amylases with soluble starch, granular starch, and maltooligosaccharides, 534
Kobrehel, K. *See* J. H. Wong, 113
Köksel, H., F. Şahbaz, and Ö. Özboy. Influence of wheat-drying temperatures on the birefringence and X-ray diffraction patterns of wet-harvested wheat starch, 481
Kozub, G. C. *See* M. S. Kaldy, 77
Kruger, J. E., and D. W. Hatcher. NOTE: A sensitive kinetic microassay for the determination of wheat α-amylase, 234
———. *See* D. W. Hatcher, 51
Kruk, M. *See* E. Souza, 280
Kumar, A., and B. M. Chauhan. Effects of phytic acid on protein digestibility (in vitro) and HCl-extractability of minerals in pearl millet sprouts, 504

Lafiandra, D. *See* S. Masci, 581
Lagimonière, M. *See* P. Gélinas, 219
Laksanalamai, V., and S. Ilangantileke. Comparison of aroma compound (2-acetyl-1-pyrroline) in leaves from pandan (*Pandanus amaryllifolius*) and Thai fragrant rice (Khao Dawk Mali-105), 381
Langemeier, J. *See* D. E. Rogers, 558
Lanning, S. P. *See* M. E. McLendon, 607
Larsen, D. M., C. S. Setser, and J. M. Faubion. Effects of flour type and dough retardation time on the senory characteristics of pizza crust, 647
Laskawy, G. *See* R. Sarwin, 553
Law, R. *See* W. R. Morrison, 385
Lawton, J. W., and Y. V. Wu. Thermal behavior of annealed acetic acid-soluble wheat gluten, 471
Lee, B. L. *See* C. Y. Lii, 188
Lee, C. C., J. A. Love, and L. A. Johnson. Sensory and physical properties of cakes with bovine plasma products substituted for egg, 18
Lee, Y. C. *See* K. O. Kim, 440
Leffler, R. G. *See* W. R. Windham, 662
Levesque, J. *See* J.-P. Rey, 759
Lew, E. J.-L. *See* S. Masci, 581
Li, X. *See* A. P. Resurreccion, 101
Lii, C. Y., and B. L. Lee. Heating A-, B-, and C-type starches in aqueous sodium chloride: Effects of sodium chloride concentration and moisture content on differential scanning calorimetry thermograms, 188
Lim, S., and P. A. Seib. Preparation and pasting properties of wheat and corn starch phosphates, 137
———, and ———. Location of phosphate esters in a wheat starch phosphate by ^{31}P-nuclear magnetic resonance spectroscopy, 145
Lin, W. D. A., G. Lookhart, and R. C. Hoseney. Partially purified proteolytic enzymes from wheat flour and their effect on elongational viscosity of cracker sponges, 448
———. *See* D. E. Rogers, 183
Lineback, D. R. *See* S. S. Heddleson, 744
Lingnert, H. *See* L. H. Dimberg, 637
Litchfield, J. B. *See* A. H. Mistry, 360
Liu, K., R. D. Phillips, and K. H. McWatters. Induced hard-to-cook state in cowpeas by freeze-thawing and calcium chloride soaking, 193
Lookhart, G. L., T. S. Cox, and O. K. Chung. Statistical analyses of gliadin reversed-phase high-performance liquid chromatography patterns of hard red spring and hard red winter wheat cultivars grown in a common environment: Classification indices, 430
———. *See* W. D. A. Lin, 448
———. *See* D. E. Rogers, 183
———. *See* K. A. Tilley, 602
Lorenz, K. J. *See* K. M. Dahlin, 329
———. *See* G. S. Ranhotra, 303
Love, J. A. *See* C. C. Lee, 18
Lukow, O. M. *See* A. Hussain, 483
Lusas, E. W. *See* H. D. Almeida-Dominguez, 214

Malcolmson, L. J., and R. R. Matsuo. Effects of cooking water composition on stickiness and cooking loss of spaghetti, 272
———, and ———. Effects of dry temperature and farina blending on spaghetti quality using response surface methodology, 1
———, ———, and R. Balshaw. Textural optimization of spaghetti using response surface methodology: Effects of drying temperature and durum protein level, 417
Malouf, R. B. *See* D. E. Rogers, 558
Mameesh, M. S., and M. Tomar. Phytate content of some popular Kuwaiti foods, 502
Mannheim, A., and M. Cheryan. Water-soluble zein by enzymatic modification in organic solvents, 115
Mariani, B. M. *See* M. G. D'Egidio, 67
———. *See* P. Novaro, 716
Marks, D. P., R. L. Stroshine, and L. D. Dunkle. Effects of storage history and hybrid on carbon dioxide production by rewetted shelled corn, 656
Marquardt, R. R. *See* D. Boros, 575
Marshall, W. E., J. I. Wadsworth, L. R. Verma, and L. Velupillai. Determining the degree of gelatinization in parboiled rice: Comparison of a subjective and an objective method, 226
Martin, J. M. *See* M. E. McLendon, 607
Marx, D. B. *See* N. Sharma, 63
Masci, S., D. Lafiandra, E. Porceddu, E. J.-L. Lew, H. P. Tao, and D. D. Kasarda. D-Glutenin subunits: N-Terminal sequences and evidence for the presence of cysteine, 581
Matsumoto, T. *See* Y. Hibi, 671
Matsuo, R. R. *See* L. J. Malcolmson, 1, 272, 417
Matt, J. *See* S. L. Edward, 748
Mauromoustakos, A. *See* K. E. Bennett, 734
Mawhinney, T. P. *See* K. A. Tilley, 602
Mazhar, H., and A. Chandrashekar. Differences in kafirin composition during endosperm development and germination in sorghum cultivars of varying hardness, 667
McCleary, B. V. *See* T. S. Gibson, 47
McDonald, C. E. *See* M. Sarwar, 405
McDonough, C. M. *See* R. Young, 179
McGuire, C. F. *See* M. E. McLendon, 607
McLendon, M. E., S. P. Lanning, C. F. McGuire, J. M. Martin, and L. E. Talbert. Variation of seed storage proteins within hard red spring wheat cultivars and effect on end-use properties, 607
McWatters, K. H. *See* K. Liu, 193
Mettler, E., and W. Seibel. Effects of emulsifiers and hydrocolloids on whole wheat bread quality: A response surface methodology study, 373
Miller, R. A., and R. C. Hoseney. The role of xanthan gum in white layer cakes, 585
Miller, S. S., P. J. Wood, L. N. Pietrzak, and R. G. Fulcher. Mixed linkage β-glucan, protein content, and kernel weight in *Avena* species, 231
Mistry, A. H., X. Wu, S. R. Eckhoff, and J. B. Litchfield. NOTE: Wet-milling characteristics of high-temperature, high-humidity maize, 360
Mohamed, A. A., B. R. Hamaker, and A. Aboubacar. Effects of flour-to-water ratio and time of testing on sorghum porridge firmness as determined by a uniaxial compression test, 739
Monsalve-Gonzalez, A., and Y. Pomeranz. Effect of spring and winter growth habitat on compositional, milling, and baking characteristics of winter wheats, 354
Moore, K. J. *See* R. Graybosch, 95
Moreau, D. *See* X. Rouau, 626
Morrison, W. R., R. F. Tester, C. E. Snape, R. Law, and M. J. Gidley. Swelling and gelatinization of cereal starches. IV. Some effects of lipid-complexed amylose and free amylose in waxy and normal barley starches, 385

Nagao, S. *See* K. Shiiba, 209
Nardi, S. *See* M. G. D'Egidio, 67, 298
———. *See* P. Novaro, 716
Neumann, P. E. *See* G. H. Ryu, 291
Newman, C. W. *See* L. Wang, 499
Newman, R. K. *See* L. Wang, 499
Neyreneuf, O., and B. Delpuech. NOTE: Freezing experiments on yeasted dough slabs. Effects of cryogenic temperatures on the baking performance, 109
Nimbona, C. *See* J. H. Wong, 113
Norris, K. H. *See* S. R. Delwiche, 29
Novaro, P., M. G. D'Egidio, B. M. Mariani, and S. Nardi. Combined effect of protein content and high-temperature drying systems on pasta cooking quality, 716
———. *See* M. G. D'Egidio, 67
Nys, M. *See* I. DeFloor, 526
———. *See* S. P. Roels, 318
———. *See* S. Vanhamel, 306

Oda, T. *See* S. Aoe, 423
Okada, K. *See* K. Shiiba, 209
Okita, T. W. *See* A. P. Resurreccion, 101
Özboy, Ö. *See* H. Köksel, 481

Pan, Z. *See* S. R. Eckhoff, 723
Paulis, J. W. *See* F. C. Felker, 685
Perez, C. M., C. P. Villareal, B. O. Juliano, and C. G. Biliaderis. Amylopectin-staling of cooked nonwaxy milled rices and starch gels, 567
———. *See* C. G. Biliaderis, 512
———. *See* B. O. Eggum, 275
———. *See* B. O. Juliano, 613, 650
Peterson, C. J. *See* R. Graybosch, 95, 458
Peterson, D. M., and A. A. Qureshi. Genotype and environment effects on tocols of barley and oats, 157, *618*
Phillips, R. D. *See* K. Liu, 193
Pietrzak, L. N. *See* S. S. Miller, 231
Pollak, L. *See* Y.-J. Wang, 171, 199, 521
Pomeranz, Y. *See* A. D. Bettge, 36
———. *See* Z. Czuchajowska, 701
———. *See* L. Gruchala, 163
———. *See* A. Monsalve-Gonzalez, 354
Ponte, J. G., Jr. *See* W.-M. Wang, 707
Porceddu, E. *See* S. Masci, 581
Potnis, P. S. *See* G. S. Ranhotra, 364
Pousset, J.-L. *See* J.-P. Rey, 759

Qureshi, A. A. *See* D. M. Peterson, 157, *618*

Raczyńska-Bojanowska, K. *See* K. Rybka, 55
Ramírez-Wong, B., V. E. Sweat, P. I. Torres, and L. W. Rooney. Development of two instrumental methods for corn masa texture evaluation, 286
———. *See* P. I. Torres, 8
Ranhotra, G. S., J. A. Gelroth, and G. J. Eisenbraun. NOTE: Correlation between Chopin and AACC methods of determining damaged starch, 236
———, ———, and B. K. Glaser. Effect of pectin-wheat bran blends on rat blood lipid and fecal responses and on muffin quality, 530
———, ———, ———, K. J. Lorenz, and D. L. Johnson. Composition and protein nutritional quality of quinoa, 303
———, ———, ———, and P. S. Potnis. NOTE: Lipidemic responses in rats fed flaxseed oil and meal, 364
———. *See* D. E. Rogers, 558
Rausch, K. D., S. K. Singh, and S. R. Eckhoff. NOTE: Magnesium bisulfite and sodium bisulfite as alternative steeping agents for wet milling, 489
———. *See* S. R. Eckhoff, 723
Reilly, P. J. *See* M. K. Dowd, 204
Resurreccion, A. P., X. Li, T. W. Okita, and B. O. Juliano. Characterization of poorly digested protein of cooked rice protein bodies, 101
Rey, J.-P., J.-L. Pousset, J. Levesque, and P. Wanty. NOTE: Isolation and composition of a natural dye from the stems of *Sorghum bicolor* (L.) Moench subsp. *americanum caudatum*, 759
Reynaert, H. R. *See* C. J. A. Vinkx, 311
Rielly, C. D. *See* G. M. Campbell, 517
Rodis, P., L.-F. Wen, and B. P. Wasserman. Assessment of extrusion-induced starch fragmentation by gel-permeation chromatography and methylation analysis, 152
Roels, S. P., G. Cleemput, X. Vandewalle, M. Nys, and J. A. Delcour. Bread volume potential of variable-quality flours with constant protein level as determined by factors governing mixing time and baking absorption levels, 318
———. *See* G. Cleemput, 324
Rogers, D. E., R. C. Hoseney, G. L. Lookhart, S. P. Curran, W. D. A. Lin, and R. G. Sears. Milling and cookie baking quality of near-isogenic lines of wheat differing in kernel hardness, 183
———, R. B. Malouf, J. Langemeier, J. A. Gelroth, and G. S. Ranhotra. Stability and nutrient contribution of β-carotene added to selected bakery products, 558
Ronalds, J. A. *See* M. L. Bason, 269
Rooney, L. W. *See* H. D. Almeida-Dominguez, 14, 214
———. *See* C. P. Friend, 252
———. *See* B. Ramirez-Wong, 286
———. *See* S. O. Serna-Saldivar, 762
———. *See* E. L. Suhendro, 412
———. *See* P. I. Torres, 8
———. *See* R. Young, 179
Rouau, X., and D. Moreau. Modification of some physicochemical properties of wheat flour pentosans by an enzyme complex recommended for baking, 626
Rybka, K., J. Sitarski, and K. Raczyńska-Bojanowska. Ferulic acid in rye and wheat grain and grain dietary fiber, 55
Ryu, G. H., P. E. Neumann, and C. E. Walker. Effects of some baking ingredients on physical and structural properties of wheat flour extrudates, 291

Sadd, P. A. *See* G. M. Campbell, 517
Şahbaz, F. *See* H. Köksel, 481
Sarwar, M., and C. E. McDonald. Detection of bread wheat farina adulterant in durum wheat semolina and pasta dried at low, high, and ultra-high temperatures, 405
Sarwin, R., G. Laskawy, and W. Grosch. Changes in the levels of glutathione and cysteine during the mixing of doughs with L-*threo*- and D-*erythro*-ascorbic acid, 553
Schatzki, T. F. *See* P. M. Keagy, 696
Schwarz, P. B. *See* G. Yang, 589
Sears, R. G. *See* D. E. Rogers, 183
Seguchi, M. NOTE: Effect of wheat aging on starch-granule surface proteins, 362
Seib, P. A. *See* W. S. Kim, 367
———. *See* S. Lim, 137, 145
———. *See* A. Xu, 463
Seibel, W. *See* E. Mettler, 373
Sempé, M. E. *See* L. E. Campaña, 760
Seo, Y. W. *See* R. Graybosch, 458
Serna-Saldivar, S. O. , M. H. Gomez, H. D. Almeida-Dominguez, A. Islas-Rubio, and L. W. Rooney. NOTE: A method to evaluate the lime-cooking properties of corn (*Zea mays*), 762
———. *See* H. D. Almeida-Dominguez, 14, 214
———. *See* P. I. Torres, 8
Setser, C. S. *See* D. M. Larsen, 647
Sharma, N., M. A. Hanna, and Y. R. Chen. Flow behavior of wheat flour-water dough using a capillary rheometer. I. Effect of capillary geometry, 59
———, and D. B. Marx. Flow behavior of wheat flour-water dough using a capillary rheometer. II. Effects of water, protein, mix, and rest time, 63
Shelton, D. R. *See* L. A. Grant, 676
Shibli, R. *See* K. I. Ereifej, 486
Shiiba, K., H. Yamada, H. Hara, K. Okada, and S. Nagao. Purification and characterization of two arabinoxylans from wheat bran, 209
Shull, J. M. *See* J. E. Fannon, 611
———. *See* J. J. Watterson, 452
Siebenmorgen, T. J. *See* K. E. Bennett, 734
———. *See* H. Sun, 727
Sievert, D., and P. Würsch. Thermal behavior of potato amylose and enzyme-resistant starch from maize, 333
Siffring, K., and B. L. Bruinsma. Effects of proof temperature on the quality of pan bread, 351
Singh, S. K. *See* K. D. Rausch, 489
Sitarski, J. *See* K. Rybka, 55
Skerritt, J. H. *See* J. L. Andrews, 241
———. *See* S. L. Edward, 748
Slominski, B. A. *See* D. Boros, 575
Snape, C. E. *See* W. R. Morrison, 385
Sokhey, A. S., and R. Chinnaswamy. Chemical and molecular properties of irradiated starch exudates, 260
Song, A. *See* P. J. Wu, 443
Sotelo, A. *See* L. Cornejo, 572
Souza, E., J. M. Tyler, K. D. Kephart, and M. Kruk. Genetic improvement in milling and baking quality of hard red spring wheat cultivars, 280
Spillman, C. K. *See* S. R. Eckhoff, 257
Stauffer, C. E. NOTE: Trypsin inhibitor measurement: Effect of order of reagent addition, 111
Stearns, M. *See* R. Graybosch, 95
Stroshine, R. L. *See* B. P. Marks, 656
Suhendro, E. L., R. D. Waniska, and L. W. Rooney. Effects of added proteins in wheat tortillas, 412
Sun, H., and T. J. Siebenmorgen. Milling characteristics of various rough rice kernel thickness fractions, 727
Sweat, V. E. *See* B. Ramirez-Wong, 286
Symons, S. J., and J. E. Dexter. Relationship of flour aleurone fluorescence to flour refinement for some Canadian hard common wheat classes, 90

Talbert, L. E. *See* M. E. McLendon, 607
Tao, H. P. *See* S. Masci, 581
Tatsumi, K. *See* S. Aoe, 423

Tester, R. F. *See* W. R. Morrison, 385
Testin, R. F. *See* A. Gennadios, 426
Thompson, D. B. *See* R. C. Yuan, 81
——. *See* G. R. Ziegler, 247
Tilley, K. A., G. L. Lookhart, R. C. Hoseney, and T. P. Mawhinney. Evidence for glycosylation of the high molecular weight glutenin subunits 2, 7, 8, and 12 from Chinese spring and TAM 105 wheats, 602
Tomar, M. *See* M. S. Mameesh, 502
Tonogai, J. R. *See* C. G. Biliaderis, 512
Torres, P. I., B. Ramírez-Wong, S. O. Serna-Saldivar, and L. W. Rooney. Effect of sorghum flour addition on the characteristics of wheat flour tortillas, 8
——. *See* B. Ramírez-Wong, 286
Trahanovsky, W. S. *See* M. K. Dowd, 204
Tso, C. C. *See* S. R. Eckhoff, 723
Tyler, J. M. *See* E. Souza, 280

Vallega, V. *See* M. G. D'Egidio, 298
Vandewalle, X. *See* S. P. Roels, 318
Vanhamel, S., G. Cleemput, J. A. Delcour, M. Nys, and P. L. Darius. Physicochemical and functional properties of rye nonstarch polysaccharides. IV. The effect of high molecular weight water-soluble pentosans on wheat-bread quality in a straight-dough procedure, 306
Van Oort, M. *See* G. Cleemput, 324
Velupillai, L. *See* W. E. Marshall, 226
Verma, L. R. *See* W. E. Marshall, 226
Vick, B. A. *See* G. Yang, 589
Villareal, C. P., B. O. Juliano, and S. Hizukuri. Varietal differences in amylopectin staling of cooked waxy milled rices, 753
——. *See* C. M. Perez, 567
Vinkx, C. J. A., H. R. Reynaert, P. J. Grobet, and J. A. Delcour. Physicochemical and functional properties of rye nonstarch polysaccharides. V. Variability in the structure of water-soluble arabinoxylans, 311

Wadsworth, J. I. *See* W. E. Marshall, 226
Walker, C. E. *See* G. H. Ryu, 291
Wang, L., Q. Xue, R. K. Newman, and C. W. Newman. Enrichment of tocopherols, tocotrienols, and oil in barley fractions by milling and pearling, 499
Wang, W.-J., P. White, L. Pollak, and J. Jane. Characterization of starch structures of 17 maize endosperm mutant genotypes with Oh43 inbred line background, 171
Wang, W.-M., and C. F. Klopfenstein. Effect of twin-screw extrusion on the nutritional quality of wheat, barley, and oats, 712
——, ——, and J. G. Ponte, Jr. Effects of twin-screw extrusion on the physical properties of dietary fiber and other components of whole bread and wheat bran and on the baking quality of the wheat bran, 707
Wang, Y.-J., P. White, and L. Pollak. Physicochemical properties of starches from mutant genotypes of the Oh43 inbred line, 199
——, ——, ——, and J. Jane. Amylopectin and intermediate materials in starches from mutant genotypes of the Oh43 inbred line, 521
Waniska, R. D. *See* H. D. Almeida-Dominguez, 214
——. *See* C. P. Friend, 252
——. *See* E. L. Suhendro, 412
——. *See* R. Young, 179
Wannerberger, L., and A.-C. Eliasson. Differential scanning calorimetry studies on rye flour-milling streams, 196
Wanty, P. *See* J.-P. Rey, 759
Wasserman, B. P. *See* K.-Y. Chan, 22, 27
——. *See* P. Rodis, 152
Watterson, J. J., J. M. Shull, and A. W. Kirleis. Quantitation of α-, β-, and γ-kafirins in vitreous and opaque endosperm of *Sorghum bicolor*, 452
Wehling, R. L., D. S. Jackson, D. G. Hooper, and A. R. Ghaedian. Prediction of wet-milling starch yield from corn by near-infrared spectroscopy, 720
Weller, C. L. *See* A. Gennadios, 426
Wen, L.-F. *See* P. Rodis, 152
White, P. J. *See* K. Cho, 539
——. *See* J. J. Huang, 42
——. *See* Y.-J. Wang, 171, 199, 521
Wilson, J. D. *See* D. B. Bechtel, 238
Windham, W. R., C. S. Gaines, and R. G. Leffler. Effect of wheat moisture content on hardness scores determined by near-infrared reflectance and on hardness score standardization, 662
Wong, J. H., K. Kobrehel, C. Nimbona, B. C. Yee, A. Balogh, F. Kiss, and B. B. Buchanan. NOTE: Thioredoxin and bread wheat, 113
Wood, P. J. *See* S. S. Miller, 231
Wrigley, C. W. *See* M. L. Bason, 269
Wu, P. J., A. Song, and S. R. Eckhoff. Enzyme treatment of popcorn ears to facilitate shelling, 443
Wu, X. *See* S. R. Eckhoff, 723
——. *See* A. H. Mistry, 360
Wu, Y. V. *See* J. W. Lawton, 471
Würsch, P. *See* D. Sievert, 333

Xu, A., and P. A. Seib. Structure of tapioca pearls compared to starch noodles from mung beans, 463
Xue, Q. *See* L. Wang, 499

Yamada, H. *See* K. Shiiba, 209
Yamauchi, M. *See* S. Aoe, 423
Yang, G., P. B. Schwarz, and B. A. Vick. Purification and characterization of lipoxygenase isoenzymes in germinating barley, 589
Yee, B. C. *See* J. H. Wong, 113
Yeh, A.-I., and S.-L. Yeh. Some characteristics of hydroxypropylated and cross-linked rice starch, 596
Yeh, S.-L. *See* A.-I. Yeh, 596
Young, R., M. H. Gomez, C. M. McDonough, R. D. Waniska, and L. W. Rooney. Changes in sorghum starch during parboiling, 179
Yuan, R. C., D. B. Thompson, and C. D. Boyer. Fine structure of amylopectin in relation to gelatinization and retrogradation behavior of maize starches from three *wx*-containing genotypes in two inbred lines, 81

Zayas, I. *See* D. B. Bechtel, 238
Zhang, X., and G. H. Brusewitz. Water absorption by cracked mustard, 133
Ziegler, G. R., D. B. Thompson, and J. Casasnovas. Dynamic measurement of starch granule swelling during gelatinization, 247

Subject Index

Page number of erratum is in italic.

Acknowledgment of reviewers, vi
Additives, emulsifiers and hydrocolloids and whole wheat quality (Mettler and Seibel), 373
Adhesion, application of Dahlquist criterion to dough (Heddleson et al), 744
Albumin fraction, associated carbohydrate of spring, winter, and soft wheats (Figueroa and Khan), 689
Alkaline, cooking of corn (Serna-Saldivar et al), 762
Amino acid, sequencing; N-terminal sequences of D-glutenin subunits (Masci et al), 581
α-Amylase
—induction and secretions in wheat aleurones and half-seeds (Bernier and Ballance), 127
—reaction patterns of maize, with starch and oligosaccharides (Knutson), 534
—sensitive microplate assay for measurement of (Kruger and Hatcher), 234
—side activities, effects on isolated pentosans (Rouau and Moreau), 626
Amylopectin
—from *wx*-containing genotypes (Yuan et al), 81
—staling of cooked nonwaxy milled rice and starch gels (Perez et al), 567
—staling in cooked waxy milled rice (Villareal et al), 753
—in starches from mutant corn genotypes (Wang et al), 521
Amylose
—content; effect on sorghum porridge firmness (Mohamet et al), 739
—determination in spaghetti and semolina (Grant et al), 676
—extender mutant of IR36 rice (Eggum et al), 275
—lipid-complexed and lipid-free in barley starch (Morrison et al), 385
—potato thermal behavior from maize (Sievert and Würsch), 333
Antibody, dough-strength prediction in early generation screening (Andrews et al), 241
Antioxidants, new type found in oats (Dimberg et al), 637
Arabinoxylan
—from wheat bran; purification and characterization of (Shiiba et al), 209
—heterogeneity in the structure of water-soluble pentosans from wheat (Cleemput et al), 324
—rye, variations in structure (Vinkx et al), 311
—solubilization by pentosanases, viscosity in solution (Rouau and Moreau), 626
—water-soluble, extract viscosity prepared from rye as indirect assay for content of (Boros et al), 575
—wheat endosperm, structural heterogeneity (Izydorczyk and Biliaderis), 641
Aroma compound, Thai fragrant rice compared in pandan leaves (Laksanalamai and Ilangantileke), 381
Ascorbic acid, mechanism of flour improvement (Sarwin et al), 553

Bagels, stability of β-carotene in (Rogers et al), 558
Bakery products, stability of β-carotene in (Rogers et al), 558
Baking
—absorption and mixing time and bread volume (Roels et al), 318
—bread, proof time and temperature effect on volume (Siffring and Bruinsma), 351
—extrusion effects on the functionality of wheat bran in whole wheat bread (Wang et al), 707
—freezing experiments on yeasted dough slabs, cryogenic temperature effects on (Neyreneuf and Delpuech), 109
—of HRS wheat cultivars, genetic improvement (Souza et al), 280
—influence of gluten components and flour lipids on cake volume and cookie diameter in soft white wheat (Kaldy et al), 77
—ingredient effect (Ryu et al), 291
Barley
—bran; extraction and enrichment of β-glucan from (Bhatty), 73
—cholesterol-lowering effects in hamsters of GEB and dehulled whole (Kahlon et al), 435
—enrichment of tocopherols, tocotrienols, and oil in, by milling and pearling (Wang et al), 499
—hull-less, physicochemical properties of bran (Bhatty), 397
—lipoxygenase isoenzymes in (Yang et al), 589
—malting, sprout damage test (Bason et al), 269
—starches; lipid-complexed and lipid-free amylose in waxy and nonwaxy (Morrison et al), 385
—tocols in (Peterson and Qureshi), 157, *618*
Bran
—barley, physicochemical properties of (Bhatty), 397
—barley and oat, extraction and enrichment of β-glucan from (Bhatty), 73
Bread
—volume; mixing time and baking absorption effect (Roels et al), 318
—wheat; thioredoxin and (Wong et al), 113
Bread and breadmaking
—effect of wheat starch, cassava starch, and cassava flour on breadmaking potential of wheat flour (DeFloor et al), 526
—optimum formulation for whole wheat bread (Mettler and Seibel), 373
—prediction using multivariate statistics (Graybosch et al), 95
—wheatless, impact of milling procedure on breadmaking potential of cassava flour (DeFloor and Delcour), 616
Brown rice, ethanol containing antioxidant or chelator to produce (Champagne and Hron), 562

Cake
—effect of xanthan gum (Miller and Hoseney), 585
—egg substituted for bovine plasma products (Lee et al), 18
—stability of β-carotene in (Rogers et al), 558
Calcium, chloride soaking; for induced hard-to-cook state in cowpeas (Liu et al), 193
Calorimetry, measurement of thermal properties (Biliaderis et al), 512
Carbohydrate
—of albumin fraction of spring, winter, and soft wheats (Figueroa and Khan), 689
—in vitro digestibility of lab-extruded cereal grains (Dahlin and Lorenz), 329
Cereal, and chicken mixtures; for infantile lactose intolerance (Cornejo et al), 572
Cereal grains, in vitro carbohydrate digestibility after laboratory extrusion (Dahlin and Lorenz), 329
Cholesterol
—effect of defatted rice bran, soluble dietary fiber (Aoe et al), 423
—lowering effects in hamsters of GEB, dehulled whole barley, rice bran, and oat bran (Kahlon et al), 435
Chromatography
—of albumin fraction of spring, winter, and soft wheats (Figueroa and Khan), 689
—preparative size-exclusion of gliadins (Huebner and Bietz), 506
—of two arabinoxylans from wheat bran (Shiiba et al), 209
Cookies
—baking; effect of wheat hardness of near-isogenic lines (Rogers et al), 183
—stability of β-carotene in (Rogers et al), 558
Corn (see also Maize)
—alternative steeping agents for wet milling (Rausch et al), 489
—amylopectin and intermediate materials in mutant starches (Wang et al), 521
—effects of storage history and hybrid on CO_2 production by (Marks et al), 656
—enzyme modification and ultrafiltration of zein (Mannheim and Cheryan), 115
—flour and chicken mixtures; evaluation of nutritive qualities (Cornejo et al), 572
—kernel vitreosity determination by video image analysis (Felker and Paulis), 685
—low molecular weight organic composition of ethanol stillage from (Dowd et al), 204
—for making masa for tortillas (Ramirez-Wong et al), 286
—physicochemical properties of starches from mutant genotypes (Wang et al), 199
—popcorn, enzyme treatment to facilitate shelling (Wu et al), 443
—prediction of starch yield by NIR spectroscopy (Wehling et al), 720
—soft-endosperm, high-lysine, short steep times for wet milling (Fox and Eckhoff), 402
—starch; preparation of phosphate ester (Lim and Seib), 137
—starch granule swelling; dynamic measurement of (Ziegler et al), 247
—starch structure in 17 mutant genotypes (Wang et al), 171
—wet-milling to increase reproducibility and accuracy of product yields (Eckhoff et al), 723
Corn meal
—extrusion effect on thiol and disulfide bond content (Chan and Wasserman), 22

—protein assay of (Chan and Wasserman), 27
Corn starch
—extrusion cooking (Rodis et al), 152
—waxy, interaction with monoglycerides (Huang and White), 42
Cowpeas
—induced hard-to-cook state in, by freeze-thawing and calcium chloride soaking (Liu et al), 193
—production and nutritional value of weaning foods from (Almeida-Dominguez et al), 14
Crackers, isolation of proteolytic enzyme that modifies elongational viscosity (Lin et al), 448
Cysteine
—detection of by fluorescent labelling (Masci et al), 581
—quantitative analysis in flours and doughs (Sarwin et al), 553

Dietary fiber
—effects of twin-screw extrusion on dietary fiber solubility (Wang et al), 707
—of wheat and rye grain, ferulic acid in (Rybka et al), 55
Differential scanning calorimetry
—for annealing acetic acid-soluble wheat gluten (Lawton and Wu), 471
—of enzyme-resistant starch (Gruchala and Pomeranz), 163
—for measurement of starch granule swelling during gelatinization (Ziegler et al), 247
—of parboiled rice, comparison with rice grading method (Marshall et al), 226
—of rye flour-milling streams (Wannerberger and Eliasson), 196
—for thermal properties of potato amylose and enzyme-resistant starch from maize (Sievert and Würsch), 333
—thermograms; NaCl and moisture effects on (Lii and Lee), 188
Disulfide, bonds; direct quantification in cereal proteins (Chan and Wasserman), 22
Dough
—density measurement and interpretation (Campbell et al), 517
—elongational viscosity, isolation of proteolytic enzyme that modifies (Lin et al), 448
—frozen, effect of yeast sampling (Gélinas et al), 219
—mixing properties; effects of charge, ADBs, and oxidants (Kerr et al), 633
—retardation time and flour type on sensory characteristics (Larsen et al), 647
—rheological criterion for adhesion, Dahlquist criterion applicability (Heddleston et al), 744
—rheological properties and pastry baking (Hay), 392
—rheology; capillary rheometer for flow behavior, capillary geometry effect (Sharma et al), 59
—rheology; capillary rheometer for flow behavior, effects of water, protein, mix, and rest time (Sharma et al), 63
—yeasted slabs; freezing experiments on, cryogenic temperature effect on baking performance (Neyreneuf and Delpuech), 109
Drying
—effect of drying temperatures on starch properties (Köksel et al), 481
—temperature and high protein content effect on pasta quality (Novaro et al), 716

Electrophoresis
—acid-PAGE and SDS-PAGE of durum wheat proteins (Hussain and Lukow), 483
—SDS-PAGE and UREA-PAGE of wheat albumin and fractions from gel filtration chromatography (Figueroa and Khan), 689
Endosperm, maize, zein composition (Dombrink-Kurtzman and Bietz), 105
Enzymes
—α-amylase, (1→3)-β-glucanase and (1→3),(1→4)-β-glucanase in wheat (Bernier and Ballance), 127
—assay for wheat α-amylase (Kruger and Hatcher), 234
—lipoxygenase isoenzymes in germinating barley (Yang et al), 589
—modification of corn proteins (Mannheim and Cheryan), 115
—polyphenol oxidase distribution in Canadian wheat flour millstreams (Hatcher and Kruger), 51
—proteolytic, isolation from cracker flour (Lin et al), 448
—reaction patterns of maize α-amylases (Knutson), 534
—resistance to amylolysis of retrograded amylose solutions (Eerlingen et al), 345
—resistance to amylolysis of wheat starch gel (Eerlingen et al), 339
Erratum, vol. 70, no. 2, p. 157, *618*
Ethanol
—containing antioxidant or chelator to produce brown rice products (Champagne and Hron), 562
—low molecular weight organic composition of stillage from distillation of (Dowd et al), 204
Extrusion
—effects on the cholesterol-lowering properties of oats, wheat, and barley (Wang and Klopfenstein), 712
—effects on in vitro carbohydrate digestibility of laboratory-extruded cereal grains (Dahlin and Lorenz), 329
—effects of physical properties of wheat and wheat bran (Wang et al), 707
—ingredient effect (Ryu et al), 291
—of starch, chemical and molecular properties of (Sokhey and Chinnaswamy), 260
—twin-screw, of rice flour with salt and sugar (Hsieh et al), 493
Extrusion cooking
—effects on starch structure (Rodis et al), 152
—pearl millet for production of weaning foods (Almeida-Dominguez et al), 214

Fiber
—in oats (Cho and White), 539
—soluble, extraction from defatted rice bran (Aoe et al), 423
Flaxseed, oil and meal; lipidemic responses (Ranhotra et al), 364
Flour
—aleurone fluorescence relation to flour refinement (Symons and Dexter), 90
—barley, physicochemical properties of (Bhatty), 397
—diploid wheat, ash, carotene, gluten, and protein content (D'Egidio et al), 298
—effect of strength on dough density (Campbell et al), 517
—fractionation into gluten and starch (Czuchajowska and Pomeranz), 701
—rheological properties of, from a landrace and durum wheat cultivars grown in Jordan (Ereifej and Shibli), 486
—from spring and winter wheats (Monsalve-Gonzalez and Pomeranz), 354
—type on sensory characteristics of pizza crust (Larsen et al), 647
Fractionation
—of D-glutenin subunits by free-flow IEF and RP-HPLC (Masci et al), 581
—LMWG subunits from monomeric gliadins (Huebner and Bietz), 506
—of preharvest sprouted wheat by air aspiration (Bettge and Pomeranz), 36
—of wheat albumin by Osborne method and by gel-filtration chromatography (Figueroa and Khan), 689

Gelatinization
—dynamic measurement of starch granule swelling during (Ziegler et al), 247
—properties of rye flour-milling streams (Wannerberger and Eliasson), 196
—starch, effects of lipid-complexed and lipid-free amylose (Morrison et al), 385
Gibberellic acid, induction and secretion of enzymes in wheat (Bernier and Ballance), 127
Glass transitions, of extrudates; processing-induced fragmentation and end-product attribute relation (Kaletunc and Breslauer), 548
Gliadin
—separation on size-exclusion column (Huebner and Bietz), 506
—statistical analyses of HPLC patterns (Lookhart et al), 430
β-Glucan
—content in *Avena* spp. (Miller et al), 231
—in oats (Cho and White), 539
Glutathione, quantitative analysis in flours and doughs (Sarwin et al), 553
Gluten
—films; physical and barrier properties (Gennadios et al), 426
—pentosan content and effect on baking quality in soft white wheat (Kaldy et al), 77
—thermal behavior of annealed acetic acid-soluble wheat (Lawton and Wu), 471
Glutenin
—characterization of D-glutenin subunits (Masci et al), 581
—HMW-GS binding antibodies for predicting dough strength (Andrews et al), 241
—LMW and HMW subunits, relation to dough properties and pastry baking (Hay), 392
—of wheat lines with varying number of HMW subunits (Gao and Bushuk), 475
Grain
—effect of wheat moisture content on standard method hardness score standardization (Windham et al), 662

—protectants; determination (Edward et al), 748
Guidelines for the preparation and review of papers reporting sensory evaluation data, v

Hardness
—of corn kernels determined by video image analysis (Felker and Paulis), 685
—effect of wheat moisture content on standard method hardness score standardization (Windham et al), 662
—separation of hard and soft red winter wheats using isolated starch as a criterion (Bechtel et al), 238
—wheat, milling and baking quality of near-isogenic lines (Rogers et al), 183
HPLC
—analysis of β-carotene in bakery products (Rogers et al), 558
—quantitative reversed phase analysis of glutathione and cysteine (Sarwin et al), 553
—separation of stillage components by (Dowd et al), 204
HPSEC, to determine molecular size distribution of starch and debranched starch (Bradbury and Bello), 543
Hydrocolloids, effect on processing and qualities of wheat tortillas (Friend et al), 252

IEF, free-flowing, for purification of wheat storage proteins (Masci et al), 581
Image analysis
—of isolated wheat starch (Bechtel et al), 238
—of weevil damage in wheat radiographs (Keagy and Schatzki), 696
Insect damage, detection in wheat radiographs, machine recognition (Keagy and Schatzki), 696
Instructions to authors, iii
Irradiation, of starch extrudates, chemical and molecular properties of (Sokhey and Chinnaswamy), 260

Legumes, cowpeas, induced hard-to-cook state in, by freeze-thawing and calcium chloride soaking (Liu et al), 193
Lipidic response, in rats fed flaxseed oil and meal (Ranhotra et al), 364
Lipids
—blood, pectin-wheat bran blend effect on (Ranhotra et al), 530
—content of various components and effect on baking quality in soft white wheat (Kaldy et al), 77
—effects on breadmaking quality (Graybosch et al), 95
Lipoxygenase, isoenzymes in germinating barley (Yang et al), 589

Maize (see also Corn)
—amylopectins, from *wx*-containing genotypes (Yuan et al), 81
—hard and soft endosperm; zein in (Dombrink-Kurtzman and Bietz), 105
—reaction patterns of α-amylases with starch and oligosaccharides (Knutson), 534
—wet-milling characteristics of (Mistry et al), 360
—wet milling of grits (Eckhoff et al), 257
Malt, effects of malting conditions on quality characteristics of (Kim et al), 440
Methods
—Chopin and AACC for determining damaged starch (Ranhotra et al), 236
—comparison with enzymatic starch damage assay kit (Gibson et al), 47
—dough density measurement (Campbell et al), 517
—effect of wheat moisture content on standard method hardness score standardization (Windham et al), 662
—estimating corn kernel vitreosity by video image analysis (Felker and Paulis), 685
Microassay, of wheat α-amylase (Kruger and Hatcher), 234
Microwave
—energy for drying wheat, influence on physical, chemical, and baking properties (Campaña et al), 760
—parboiling; effect on starch gelatinization (Marshall et al), 226
Millet, cowpea, extrusion cooking of millet for production of weaning foods (Almeida-Dominguez et al), 214
Milling
—and baking quality; effect of wheat hardness (Rogers et al), 183
—degree measured objectively by measuring surface fat concentrations (Bennett et al), 734
—DSC studies on rye flour-milling streams (Wannerberger and Eliasson), 196
—of HRS wheat cultivars, genetic improvement (Souza et al), 280
—quality affected by rice kernel thickness (Sun and Siebenmorgen), 727
—roller and hammer milled mustard effect on water absorption (Zhang and Brusewitz), 133
Mineral, availability; effect of phytic acid in pearl millet sprouts (Kumar and Chauhan), 504
Mixing, effect of speed on dough density (Campbell et al), 517
Mixograph, combined effects of pH, ADBs, and oxidants (Kerr et al), 633
Moisture, critical content for fissures in rough rice (Juliano and Perez), 613
Monoglycerides, interaction with waxy corn starch (Huang and White), 42
Muffins, quality affected by pectin-wheat bran blends (Ranhotra et al), 530
Mustard, water absorption of cracked seed (Zhang and Brusewitz), 133

NIR spectroscopy
—effect of wheat moisture content on standard method hardness score standardization (Windham et al), 662
—hard red wheat classified by (Delwiche and Norris), 29
—prediction of wet-milling starch yield from corn (Wehling et al), 720

Oat bran
—cholesterol-lowering effects of in hamsters (Kahlon et al), 435
—extraction and enrichment of β-glucan from (Bhatty), 73
Oats
—β-glucan content in different oat genotypes (Cho and White), 539
—β-glucan, protein content, and kernel weight in *Avena* spp. (Miller et al), 231
—identification of new phenolic antioxidants (Dimberg et al), 637
—tocols in (Peterson and Qureshi), 157, *618*
Oil, enrichment of in barley fractions by milling and pearling (Wang et al), 499

Parboiling, comparison of Instron firmness with dynamic viscoelastic properties (Edwards et al), 122
Pasta
—cooking quality evaluation by sensory judgment, total organic matter, and viscoelastograph measures at low and high drying temperatures (D'Egidio et al), 67
—detecting adulteration by analysis for sterol palmitates (Sarwar and McDonald), 405
—effect of cooking water composition on stickiness and cooking loss (Malcolmson and Matsuo), 272
—effects of drying temperature and farina blending on quality (Malcolmson et al), 1
—effects of drying temperature and protein level (Malcolmson et al), 417
—high-temperature drying and grain protein content effect on quality (Novaro et al), 716
—noodles, restriction of starch swelling in, by lipids in wheat flours (Kim and Seib), 367
Pastry, effect of flour quality characteristics and dough rheological properties on baking quality (Hay), 392
Pearl millet
—extrusion cooking for production of weaning foods (Almeida-Dominguez et al), 214
—and nutritional value of weaning foods from (Almeida-Dominguez et al), 14
—phytic acid effect on protein digestibility and mineral availability (Kumar and Chauhan), 504
Pentosans
—effects on breadmaking quality (Graybosch et al), 95
—in gluten from soft white wheat, influence on baking quality (Kaldy et al), 77
—heterogeneity in the structure of water-soluble arabinoxylans from wheat (Cleemput et al), 324
—modification of properties by enzymatic hydrolysis (Rouau and Moreau), 626
—rye, effects on mixing times, baking absorption, and bread volume (Vanhamel et al), 306
—rye, variations in structure (Vinkx et al), 311
—water-soluble, extract viscosity prepared from rye as indirect assay for content of (Boros et al), 575
Phenolics, antioxidants in oats (Dimberg et al), 637
Phytate, popular Kuwaiti food content and reduction during cooking of rice and legumes (Mameesh and Tomar), 502
Phytic acid, effect on protein digestibility and mineral availability in pearl millet sprouts (Kumar and Chauhan), 504
Polyphenol oxidase, distribution in Canadian wheat flour millstreams (Hatcher and Kruger), 51
Popcorn, enzyme treatment to facilitate shelling (Wu et al), 443

Porridge, thick, from sorghum (Mohamet et al), 739
Protein
—albumin from spring, winter, and soft wheats (Figueroa and Khan), 689
—bodies; cooked rice, poorly digested (Resurreccion et al), 101
—cereal, solubilized and particulate (Chan and Wasserman), 22
—combined effect with high-temperature drying on pasta cooking quality (Novaro et al), 716
—concentrates and prime starch from wheat flours (Czuchajowska and Pomeranz), 701
—content in *Avena* spp. (Miller et al), 231
—digestibility; effect of phytic acid in pearl millet sprouts (Kumar and Chauhan), 504
—direct colorimetric assay (Chan and Wasserman), 27
—effect of disulfide bonds on rice starch gelatinization (Hamaker and Griffin), 377
—effect in wheat tortillas (Suhendro et al), 412
—enzyme modification and ultrafiltration of zein (Mannheim and Cheryan), 115
—high molecular weight glutenin subunit characterization of (Tilley et al), 602
—influence on durum wheat quality (Hussain and Lukow), 483
—monoclonal antibodies to rye secaling (Graybosch et al), 458
—physical, chemical, functional properties (Al-Kahtani and Abou-Arab), 619
—seed storage, in HRS wheat cultivars (McLendon et al), 607
—starch-granule surface, effect of wheat flour aging (Seguchi), 362
—utilization in mutant milled rice (Eggum et al), 275
—wheat flour, separation by SE-HPLC and relation to quality (Graybosch et al), 95
—zein in endosperm, of maize (Dombrink-Kurtzman and Bietz), 105

Quinoa, composition and protein nutritional quality of (Ranhotra et al), 303

Recently accepted manuscripts, 490, 618, 758
Response surface methodology
—effect of drying temperature and farina blending on spaghetti quality (Malcolmson et al), 1
—effect of drying temperature and protein level on spaghetti texture (Malcolmson et al), 417
—for malting condition effects on quality characteristics of malt (Kim et al), 440
—optimization of whole wheat bread quality (Mettler and Seibel), 373
Rheological properties
—doughs, effect of charge, ADBs, and oxidants (Kerr et al), 633
—of flour from a landrace and durum wheat cultivars grown in Jordan (Ereifej and Shibli), 486
—maturograph and oven rise recorder, registration of dough behavior and oven rise during fermentation and baking (Mettler and Seibel), 373
Rheology
—dough, capillary rheometer for flow behavior, capillary geometry effect (Sharma et al), 59
—dough, capillary rheometer for flow behavior, effects of water, protein, mix, and rest time (Sharma et al), 63
—Instron firmness and oscillatory measurements on cooked pasta (Edwards et al), 122
—properties of diploid, durum, and common wheat flours (D'Egidio et al), 298
—uniaxial compression test for porridge firmness (Mohamed et al), 739
—viscoelastic properties of masa (Souza et al), 280
Rice
—cooked, protein bodies, poorly digested (Resurreccion et al), 101
—cooked nonwaxy milled, amylopectin staling of (Perez et al), 567
—cooked waxy milled, amylopectin staling of (Villareal et al), 753
—flour; twin-screen extrusion with salt and sugar (Hsieh et al), 493
—kernel thickness distribution (Sun and Siebenmorgen), 727
—parboiled, degree of gelatinization (Marshall et al), 226
—rough, critical moisture content for fissures in (Juliano and Perez), 613
—stable high head yield, screening for (Juliano et al), 650
—starch; effect of cross-linking and hydroxypropylation on properties of (Yeh and Yeh), 596
—starch gelatinization and paste viscosity (Hamaker and Griffin), 377
—surface fat concentration measured on samples milled in a McGill No. 2 miller at various settings (Bennet et al), 734
—Thai fragrant, aroma compound compared in pandan leaves (Laksanalamai and Ilangantileke), 381
Rice bran
—cholesterol-lowering effects of in hamsters (Kahlon et al), 435
—extraction of soluble dietary fibers (Aoe et al), 423
RP-HPLC
—detecting pasta adulteration by sterol palmitate (Sarwar and McDonald), 405
—LMW and HMW glutenins, gliadins (Hay), 392
—of LMWG subunits, both unreduced and reduced (Huebner and Bietz), 506
—of zein in endosperm, of maize (Dombrink-Kurtzman and Bietz), 105
Rye
—detection of rye secalins via immunoblotting and ELISA (Graybosch et al), 458
—DSC studies on rye flour-milling streams (Wannerberger and Eliasson), 196
—ferulic acid content in grain and dietary fiber (Rybka et al), 55
—HMW pentosan effect on wheat-bread quality in straight-dough procedure (Vanhamel et al), 306
—relationship between water-soluble pentosan content and viscosity of three extracts prepared from (Boros et al), 575
—water-soluble arabinoxylans, variations in structure (Vinkx et al), 311

Salt, for twin-screw extrusion of rice flour (Hsieh et al), 493
Scanning electron microscopy, of starch from 17 maize mutant genotypes (Wang et al), 171
SDS-PAGE
—immunoblots with antisecalin monoclonal antibody (Graybosch et al), 458
—for qualitative and quantitative differences in vitrous and opaque endosperms of sorghum (Watterson et al), 452
Sodium chloride, effects on differential scanning calorimetry thermograms (Lii and Lee), 188
Sorghum
—endosperm development and germination (Mazhar and Chandrashekar), 667
—endosperm texture and composition related to porridge quality (Mohamed et al), 739
—flour addition effect on tortillas (Torres et al), 8
—interior channels of starch granules (Fannon et al), 611
—natural dye from stems of, isolation and composition (Rey et al), 759
—starch; changes during parboiling (Young et al), 179
—vitreous and opaque endosperm, quantitation of kafirins in (Watterson et al), 452
Soybeans
—protein compared to *Moringa peregrina* (Al-Kahtani and Abou-Arab), 619
—protein films, physical and barrier properties (Gennadios et al), 426
Spaghetti, effects of drying, starch damage, sprouting on quality characteristics (Grant et al), 676
Starch
—amylopectin and intermediate materials in corn genotypes (Wang et al), 521
—amylopectins from *wx*-containing maize genotypes (Yuan et al), 81
—changes in sorghum during parboiling (Young et al), 179
—chemical and molecular properties of irradiated extrudates (Sokhey and Chinnaswamy), 260
—corn and wheat starch phosphates, preparation and properties (Lim and Seib), 137
—crystalline structure; effect of high pressure (Hibi et al), 671
—damage, Chopin and AACC methods of determining (Ranhotra et al), 236
—damage, effect on spaghetti stickiness (Grant et al), 676
—damage measured with enzymatic assay kit (Gibson et al), 47
—digestibility in mutant milled rice (Eggum et al), 275
—effect of rice protein on gelatinization (Hamaker and Griffin), 377
—enzyme-resistant from maize (Sievert and Würsch), 333
—enzyme-resistant, studies by differential scanning calorimetry (Gruchala and Pomeranz), 163
—gels; amylopectin staling of (Perez et al), 567
—granule swelling; effects of lipid-complexed and lipid-free amylose (Morrison et al), 385
—image analysis of isolated wheat starch (Bechtel et al), 238
—impact of amylose chain length on formation of enzyme-resistant, in aqueous solutions of amylose (Eerlingen et al), 345
—impact on the breadmaking potential of wheat flour (DeFloor et al), 526
—interior channels of sorghum granules (Fannon et al), 611
—molecular size distribution determined by HPSEC (Bradbury and Bello), 543
—noodles from mung beans; structure of tapioca pearls compared to (Xu and Seib), 463

—phosphate, location of ester group by ^{31}P-NMR spectroscopy (Lim and Seib), 145
—physicochemical properties from mutant genotypes (Wang et al), 199
—prime, from wheat flours (Czuchajowska and Pomeranz), 701
—reaction patterns of maize α-amylases with soluble, granular (Knutson), 534
—structure in 17 mutant genotypes (Wang et al), 171
—swelling; restriction in cooked noodles by lipids in wheat flours (Kim and Seib), 367
—time-temperature dependence of formation of enzyme-resistant, in wheat starch gel (Eerlingen et al), 339
—wheat, birefringence and X-ray diffraction patterns (Köksel et al), 481
Steeping
—alternative agents for wet milling (Rausch et al), 489
—cracked and whole mustard seed (Zhang and Brusewitz), 133
—short time, wet milling, soft-endosperm high-lysine corn (Fox and Eckhoff), 402
Sterol palmitate, detecting pasta adulteration (Sarwar and McDonald), 405
Storage, effects of history on CO_2 production by shelled corn (Marks et al), 656
Sugar, for twin-screw extrusion of rice flour (Hsieh et al), 493
Sulfhydryl, groups; direct quantification in cereal proteins (Chan and Wasserman), 22

Tapioca pearls, structure compared to starch noodles from mung bean (Xu and Seib), 463
Tests, moisture adsorption stress for screening stable high head rice yields in rough rice (Juliano et al), 650
Texture
—of cooked pasta (Edwards et al), 122
—of masa (Souza et al), 280
—optimization of spaghetti (Malcolmson et al), 417
—sorghum porridge, influence of flour-to-water ratio and time of testing (Mohamed et al), 739
—of wheat flour tortillas (Torres et al), 8
Thioredoxin, and bread wheat (Wong et al), 113
Tocols, in barley and oats (Peterson and Qureshi), 157, *618*
Tocopherols, enrichment of in barley fractions by milling and pearling (Wang et al), 499
Tocotrienols, enrichment of in barley fractions by milling and pearling (Wang et al), 499
Tortillas
—corn (Serna-Saldivar et al), 762
—wheat, effects of added proteins (Suhendro et al), 412
—wheat, effects of hydrocolloids on processing and qualities of (Friend et al), 252
—wheat flour, effect of sorghum flour addition on (Torres et al), 8
Total organic matter, for pasta cooking quality evaluation; role at low and high drying temperatures (D'Egidio et al), 67
Trypsin inhibitor, effect of order of reagent addition on measurement (Stauffer), 111

Ultrafiltration, corn proteins (Mannheim and Cheryan), 115

Viscoelastograph, measures; role for pasta cooking quality evaluation at low and high drying temperatures (D'Egidio et al), 67
Viscosity, soluble pentosans, reduction by enzymatic treatment (Rouau and Moreau), 626
Vitreosity, of corn kernels determined by video image analysis (Felker and Paulis), 685

Wet milling
—corn, alternative steeping agents for (Rausch et al), 489
—corn, to increase reproducibility and accuracy of product yields (Eckhoff et al), 723
—corn, short steep times for soft-endosperm high-lysine (Fox and Eckhoff), 402
—of high-temperature, high-humidity maize (Mistry et al), 360
—of maize grits (Eckhoff et al), 257
—prediction of starch yield from corn by NIR spectroscopy (Wehling et al), 720
Wheat
—albumin fraction from spring, winter, and soft (Figueroa and Khan), 689
—classification; spring vs. winter (Lookhart et al), 430
—composition, growth habitat, aand milling characteristics (Monsalve-Gonzalez and Pomeranz), 354
—detection of wheat-rye chromosomal translocations using monoclonal antibodies (Graybosch et al), 458
—diploid, grain, flour, and dough characteristics of (D'Egidio et al), 298
—drying with microwave energy, changes on physical, chemical, and baking properties (Campaña et al), 760
—drying temperature effect on starch properties (Köksel et al), 481
—durum, different methods in evaluating textural characteristics of cooked pasta (D'Egidio et al), 67
—effect of wheat moisture content on standard method hardness score standardization (Windham et al), 662
—endosperm arabinoxylans; structural heterogeneity (Izydorczyk and Biliaderis), 641
—ferulic acid content in grain and dietary fiber (Rybka et al), 55
—flour aging; effect on starch-granule surface proteins (Seguchi), 362
—flour aleurone fluorescence relation to flour refinement in (Symons and Dexter), 90
—flour fractionation into gluten and starch (Czuchajowska and Pomeranz), 701
—flour type; effect of on pizza crust (Larsen et al), 647
—genetic improvement in milling and baking quality of HRS (Souza et al), 280
—hardness determined from starch analysis (Bechtel et al), 238
—hard red, classification by NIR spectroscopy (Delwiche and Norris), 29
—high molecular weight glutenin subunits, evidence for glycosylation (Tilley et al), 602
—machine recognition of weevil damage in X-ray images (Keagy and Schatzki), 696
—microplate assay for wheat α-amylase (Kruger and Hatcher), 234
—omega gliadins of durum wheat (Hussain and Lukow), 483
—preharvest-sprouted, air-aspirated cleaning (Bettge and Pomeranz), 36
—proteins, lipids, and pentosans in relation to quality (Graybosch et al), 95
—quality screening using simple antibody based test (Andrews et al), 241
—seed storage protein variation within, effect on end-use properties (McLendon et al), 607
—separation of LMWG subunits from monomeric gliadins (Huebner and Bietz), 506
—starch; preparation of phosphate ester (Lim and Seib), 137
—starch phosphate, location of ester group by ^{31}P-NMR spectroscopy (Lim and Seib), 145
Wheat bran
—and pectin blends; effect on rat blood lipid and fecal responses and on muffin quality (Ranhotra et al), 530
—purification and characterization of arabinoxylans (Shiiba et al), 209

Xanthan gum, role in white layer cakes (Miller and Hoseney), 585

Yeast
—activity in frozen doughs, effects of strain, trademark, and samplings (Gélinas et al), 219
—amount necessary to proof at various temperatures (Siffring and Bruinsma), 351

Zein
—enzyme modification and ultrafiltration (Mannheim and Cheryan), 115
—in maize kernel endosperm (Dombrink-Kurtzman and Bietz), 105